

SU(3) gauge theory at finite temperature in 2 + 1 dimensions

P. Bialas,^a L. Daniel,^a A. Morel^b and B. Petersson^c

^aInstitute for Physics, Jagellonian University,
ul. Reymonta 4, 30 059 Krakow, Poland

^bService de Physique Theorique, CEA Saclay
F-91191 Gif-sur-Yvette Cedex, France

^cFakulty for Physics, University of Bielefeld,
P.O.Box 10 01 31, D-33501 Bielefeld, Germany

In this article we will discuss numerical results on screening masses and thermodynamic quantities in 2 + 1 dimensional SU(3) gauge theory. We will also compare them to perturbation theory and the dimensionally reduced model.

The SU(3) gauge theory in 2+1 dimensions is simple enough from a numerical point of view, so that it is possible with present computers to make continuum extrapolations with controlled systematic errors. Of course, there are some obvious differences between SU(3) gauge theory in 2 + 1 and 3 + 1 dimensions. In 2 + 1 dimensions the coupling constant g^2 has the dimension of a mass, and the theory is superrenormalizable. The tree level potential between heavy quarks is already logarithmically confining: $V(r) \sim \log r$. There are, however, many similarities. One may introduce a dimensionless “running” coupling constant $g_3(l)$ by the definition $g_3^2(l) \equiv lg^2$ where l is a length scale. Then $g_3^2(l) \rightarrow 0$ for $l \rightarrow 0$ and to infinity for $l \rightarrow \infty$. This is somewhat analogous to the logarithmically running coupling constant in 3 + 1 dimensional SU(3) gauge theory. In 2 + 1 dimensions the coupling constant g^2 sets the scale, and $m_i = c_i g^2$, where c_i ’s are numerical constants. From Monte Carlo simulations one knows some further properties: There is a linearly rising non-perturbative potential $V(r) \simeq \sigma_0 r$ for r large [1, 2]. There is a second order phase transition at $T_c = 0.55(1)g^2$, with the critical indices of the 2d 3states Potts model [2]. Furthermore, the glue ball masses m_{GB} are much bigger than T_c , $m_{GB} \geq 4.4T_c$ [1]. This is all qualitatively similar to 3 + 1 dimensions, where, however, the transition is weakly first order. In the gluon plasma phase $T > T_c$, one should be able to use perturbation theory. The relevant dimensionless coupling in 2 + 1 dimensions is

$$g_3^2(T) \equiv \frac{g^2}{T} \xrightarrow{T \rightarrow \infty} 0. \quad (1)$$

There are, however, infrared divergences, which are even more serious than in 3 + 1 dimensions. For the screening mass (rsp. the pressure) they appear already at order g^2 , i.e. at one (resp. two) loop(s). The infrared divergences can be tamed through resummations, e.g. through the selfconsistent perturbation theory (SCPT) introduced by




Figure 1. Screening masses from the 3D SU(3) theory on lattices with temporal extent $N_\tau = 4$ compared to the predictions of SCPT(continuous lines) and to the 2D reduced model.

D'Hoker[3]. It works in the following way. Choose the class of static gauges $\partial_0 A_0(x) = 0$ where the Polyakov loop is purely static,

$$L(\vec{x}) = Tr e^{i \frac{g}{T} A_0(\vec{x})}. \quad (2)$$

Add and subtract an explicit mass term in the static sector,

$$\frac{1}{2} m_g^2 Tr(A_0(\vec{x}))^2, \quad (3)$$

which is invariant inside this class of gauges, and perform the perturbative expansion in the theory with A_0 massive. In fact, there are no further infrared divergencies in 2+1 dimensions[3].

The second method, which is semianalytic, is the dimensional reduction from $2 + 1$ to 2 dimensions[4]. Again we choose the class of static gauges. We then integrate perturbatively over the non-static modes. This integration is by construction infrared finite. The result is an effective 2-dimensional adjoint Higgs model for the static modes with the action


$$S = \int d^2x Tr \left\{ \frac{1}{2} F_{ij}^2 + (D_i \phi)^2 - \frac{3g^2 T}{2\pi} \left[\frac{5}{2} \log 2 + 1 - \log(aT) \right] \phi^2 + \frac{g^4}{32\pi} \phi^4 \right\} \quad (4)$$

where D_i is a covariant derivative and where the coupling constants are derived from a one loop integration over the static modes. This action is systematic in $\frac{g^2}{T}$, e.g. the term ϕ^6 is multiplied by a constant proportional to $\frac{g^6}{T}$, and is neglected at high T .

The two-dimensional adjoint Higgs model has not been solved analytically. We solve it non-perturbatively by a Monte Carlo lattice simulation.

We define a screening mass for $T > T_c$ by

$$\langle Re \left(L(0) L^\dagger(\vec{x}) \right) \rangle \simeq \langle L \rangle^2 + \frac{A}{\sqrt{m|\vec{x}|}} e^{-m|\vec{x}|} \quad (5)$$


 Figure 2. Trace anomaly $(\epsilon - 2p)/T^3$, on lattices of various sizes.

In lowest order perturbation theory $m = 2m_g$ for the correlations of $ReL(\bar{x})$ and $m = 3m_g$ for the correlations of $ImL(\bar{x})$. In SCPT one has[3]

$$\frac{m_g^2}{g^2 T} = \frac{3}{2\pi} \left(\log \frac{T}{m_g} + C \right) + \mathcal{O} \left(1/\log \frac{T}{m_g} \right), \quad C = -1.0. \quad (6)$$

In Fig. 1, we show $2m_g$ compared to the screening mass in the 2 + 1 dimensional SU(3) gauge theory. We have used the formula $T/T_c = (\beta - 1.5)/3.3N_\tau$, derived from the condition $T_c/g^2 = 0.55$ for all N_τ , and the values $\beta_c(N_\tau)$ from[2]. Solving Eq. (6) where m_g/T has been replaced by $(1 - \exp(-m_g/T))$, we can get agreement only for $T/g^2 > 6$ ($T/T_c > 12$), and this only by arbitrarily choosing $C = 1.6$.

The dimensionally reduced model is in good agreement with the full theory already for $T > 1.5T_c$ as can be seen in Fig. 1. A further investigation showed that the two exchanged states in the reduced model are simple poles, not 2 gluon and 3 gluon cuts respectively[5].

For the thermodynamics we use the lattice integral method introduced in [6]. The trace anomaly, $(\epsilon - 2p)/T^3$, is presented in Fig. 2. It is qualitatively similar to the result in 3 + 1 dimensions.

Since the trace anomaly is zero for free massless gluons, we expect in perturbation theory that

$$\frac{\epsilon - 2p}{T^3} \simeq \frac{g^2}{T} f \left(\log \frac{T}{g^2} \right). \quad (7)$$

In Fig. 3 we present $(\epsilon - 2p)/T^2 g^2$ as a function of T_c/T . One observes that this quantity is slowly varying at high temperature. A comparison with perturbation theory and dimensional reduction is in progress.

In Fig. 4, we present the extrapolation of the pressure to the continuum.

This work is supported through ENRAGE (European Network on Random Geometry), a Marie Curie Research Training Network supported by the European Community's Sixth Framework Programme, network contract MRTN-CT-2004-005616.




Figure 3. Check of the scaling formula (7).




Figure 4. The pressure extrapolated to the infinite N_τ limit from the lattices of size 4×32^2 , 6×48^2 and 8×64^2 using a linear fit in $1/N_\tau^2$. The dashed line shows the Stefan-Boltzman limit.

REFERENCES

1. M. Teper, Phys. Rev **D59** (1999) 014512.
2. C. Legeland, Thesis (Bielefeld 1998), J. Engels *et al.*, Nucl. Phys. Proc. Suppl. 53 (1997) 430.
3. E. D'Hoker, Nucl. Phys. **B201** (1982) 401.
4. P. Bialas *et al.*, Nucl. Phys. **B581** (2000) 471.
5. P. Bialas *et al.*, Nucl. Phys. **B603** (2001) 369.
6. J. Engels *et al.*, Phys. Lett. **B253** (1990) 625.